arXiv:1211.6119v1 [hep-ph] 26 Nov 2012

Evading death by vacuum

A. Barroso,[1] P.M. Ferreira,[2, 1, *] I.P. Ivanov,[3, 4, †] Rui Santos,[2, 1, ‡] and João P. Silva[2, 5, §]
[1]*Centro de Física Teórica e Computacional, Faculdade de Ciências,*
Universidade de Lisboa, Av. Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal
[2]*Instituto Superior de Engenharia de Lisboa, 1959-007 Lisboa, Portugal*
[3]*IFPA, Université de Liége, Allée du 6 Août 17, bâtiment B5a, 4000 Liége, Belgium*
[4]*Sobolev Institute of Mathematics, Koptyug avenue 4, 630090, Novosibirsk, Russia*
[5]*Centro de Física Teórica de Partículas (CFTP), Instituto Superior Técnico,*
Universidade Técnica de Lisboa, 1049-001 Lisboa, Portugal
(Dated: November 28, 2012)

Quantum Mechanics and Quantum Field Theory have forced a reappraisal of the vacuum's exact nature. This is specially true in the Standard Model of particle interactions, where spontaneous symmetry breaking of the electroweak symmetry by the vacuum is mandatory for elementary particles to acquire masses. The mechanism implemented with one scalar doublet leads to a simple vacuum structure at current energies. But, the open possibility that there might be two scalar doublets, enriches the vacuum structure, allowing for the possibility that we might now be in a metastable state, which we dub the panic vacuum. Current experiments at LHC are probing the Higgs particle predicted as a result of the spontaneous symmetry breaking. Remarkably, these experiments may also be useful to dispel the panic vacuum, as we show in a simple model with two doublets and a softly broken $U(1)$ symmetry.

PACS numbers: 12.60.Fr, 14.80.Ec, 11.30.Qc, 11.30.Ly

It has been known since the inception of the quantum field theory of fundamental particles that quantum fluctuations must exist, implying that the vacuum is not "nothingness". Moreover, the gauge theory describing the electromagnetic, weak, and strong interactions (known as the Standard Model of particle physics - SM) further altered our perception of the vacuum [1]. Indeed, in the SM, fundamental particles can only obtain masses through the mechanism of spontaneous gauge symmetry breaking, achieved via the introduction of a scalar field subject to a potential with quadratic and quartic terms. The minimum of this potential is attained when the scalar field develops a non-vanishing vacuum expectation value (vev), thus imparting structure to the vacuum. This is required because all fundamental particle masses are proportional to the vev. One consequence of this mechanism is the prediction of a neutral particle, known as the Higgs scalar [2].

Over the past forty years, the SM has successfully passed numerous tests. But the mechanism of spontaneous symmetry breaking (SSB) has only recently come under experimental scrutiny, with the discovery by the LHC experiments ATLAS and CMS of a particle with properties similar to those expected for the SM Higgs [3]. The particle is produced by colliding two protons (pp) and detected, at present, mostly through its decay into two photons ($\gamma\gamma$) and its decay into two Z bosons (ZZ). It turns out that the specific production mechanism can sometimes be tagged and that it affects the detectability. In the case of $\gamma\gamma$, results are known for all production mechanisms combined, and also for production of the Higgs through the fusion of two Z or of two W bosons, known collectively as vector boson fusion (vbf).

The field content of the SM has been determined by experiment. Indeed, CP violation aside, the theory would have been viable if there were only one charged lepton, one neutrino, one up-quark, and one down-quark. But three such families have been found, with increasing masses. In the same fashion, there may be more than one scalar doublet; only experiment will tell. Of course, having (say) two scalar doublets increases the particle spectrum: there would be three neutral scalars (h and H, respectively the lightest and heaviest CP-even scalars; and A, the pseudoscalar); and a pair of conjugate charged scalars ($H^{\pm}$). But having more than one scalar doublet also enriches the vacuum structure. For example, in the SM with one scalar doublet, the potential admits only one minimum (up to the gauge transformations). In contrast, in the SM with two scalar doublets, there could coexist two neutral local minima; the global minimum with a smaller potential energy density; and a metastable state with a larger potential energy density.

This raises a troubling possibility. If we were at present in the metastable vacuum, then the scalar field could in time decay into the real vacuum, thus reheating the Universe, altering all masses, all physics, all chemistry, and invalidating our very existence. In this paper we show that even the current preliminary results can already be used to probe this question.

As an example, we consider the scalar potential of a two Higgs doublet model (2HDM) with a Peccei-Quinn symmetry given by [4, 5]

$$\begin{aligned} V_H &= m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 - m_{12}^2 \left[\Phi_1^\dagger \Phi_2 + \Phi_2^\dagger \Phi_1\right] \\ &+ \tfrac{1}{2}\lambda_1 (\Phi_1^\dagger \Phi_1)^2 + \tfrac{1}{2}\lambda_2 (\Phi_2^\dagger \Phi_2)^2 \\ &+ \lambda_3 (\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) + \lambda_4 (\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1), \end{aligned} \tag{1}$$

where all parameters are real because CP invariance is imposed. The Peccei-Quinn $U(1)$ symmetry $(\Phi_1, \Phi_2) \to (\Phi_1, e^{i\xi}\Phi_2)$ is imposed on the quartic terms and softly broken by m_{12}^2, thus guaranteeing that the pseudoscalar has a mass.

Since our Universe does not break electric charge, we know that, after SSB, only the neutral components of the scalars develop vevs, $\langle \Phi_1^0 \rangle = v_1/\sqrt{2}$ and $\langle \Phi_2^0 \rangle = v_2/\sqrt{2}$, which can be traded for $v = \sqrt{v_1^2 + v_2^2}$ and $\tan\beta = v_2/v_1$. Without loss of generality, we can take v_1 and v_2 positive and $0 \le \beta \le \pi/2$. The vevs can be obtained by minimizing V_H, leading to two coupled cubic equations. Given $m_{11}^2 \dots \lambda_4$, these can be solved numerically. Fortunately, many features can be determined without solving the equations explicitly.

In Ref. [6] one can find a proof that the potential of the most general 2HDM has at most two minima, along with conditions under which the two minima occur. Applying to our specific case, the two minima occur if and only if

$$\begin{aligned} m_{11}^2 + k^2\, m_{22}^2 &< 0, \qquad (2) \\ \sqrt[3]{x^2} + \sqrt[3]{y^2} &\le 1, \qquad (3) \end{aligned}$$

where,

$$\begin{aligned} x &= \frac{4\ k\ m_{12}^2}{m_{11}^2 + k^2\, m_{22}^2}\, \frac{\sqrt{\lambda_1 \lambda_2}}{\lambda_{34} - \sqrt{\lambda_1\lambda_2}}, \\ y &= \frac{m_{11}^2 - k^2\, m_{22}^2}{m_{11}^2 + k^2\, m_{22}^2}\, \frac{\sqrt{\lambda_1\lambda_2} + \lambda_{34}}{\sqrt{\lambda_1\lambda_2} - \lambda_{34}}, \end{aligned} \tag{4}$$

with

$$\lambda_{34} = \lambda_3 + \lambda_4 \quad \text{and} \quad k = \sqrt[4]{\frac{\lambda_1}{\lambda_2}}. \tag{5}$$

The limiting curve $\sqrt[3]{x^2} + \sqrt[3]{y^2} = 1$ is known as the astroid.

The vacuum that our Universe is currently in has $v = 2m_W/g = 246$ GeV (related to the W mass and the weak coupling g). We are particularly interested in the possibility that: i) there is a second minimum; and ii) it has vevs, v_1' and v_2', corresponding to a lower potential energy density:

$$V_H(v_1', v_2') < V_H(v_1, v_2). \tag{6}$$

For the region $\lambda_{34} > \sqrt{\lambda_1\lambda_2}$, we have developed a new discriminant

$$D = \left(m_{11}^2 - k^2 m_{22}^2\right)(\tan\beta - k). \tag{7}$$

If, in this region, $D < 0$, then Eq. (6) holds, we live in a metastable state, and we denote our current (v_1, v_2) solution the "*panic vacuum*". In the region where $\lambda_{34} < \sqrt{\lambda_1\lambda_2}$, there is no panic vacuum in this model.

Conversely, if there is only one minimum, or even if there are two minima – Eqs. (2) and (3) hold – but $D > 0$ in the region where it is defined, then (v_1, v_2) corresponds to the global minimum of the theory, and our vacuum is stable.

Problems related to ours are sometimes tackled by following the evolution of the various minima as the temperature (T) of the Universe decreased, from its inception to the present day. This involves estimates of the finite temperature contributions to the effective potential, and hinges on a variety of assumptions. Some recent work within the 2HDM can be found in Ref. [7]. It is also true that, even at $T = 0$, the nature of the vacuum changes as we increase the energy scale above 10 TeV. This occurs even in the SM, due to loop corrections to V_H, and might ultimately point to a more complete theory at high energies, encompassing the SM [8]. This is not our concern here. We are worried about a problem that affects us right now, at $T = 0$ and small energies.

What we will now show is that measurements in the laboratory can be used to reconstruct the shape of the present day ($T = 0$) tree-level scalar potential, and, thus, ascertain as to the putative presence of a lower lying minimum. Even more striking, although there are not, as of yet, enough measurements to determine all the seven $m_{11}^2 \dots \lambda_4$ parameters of V_H, the experiments at LHC are already close to ruling out the *panic vacuum* situation of Eq. (6).

If the model we are discussing describes Nature, and the three neutral scalars and conjugate pair of charged scalars are detected, then the scalar potential can be reproduced from

$$\begin{aligned} m_{12}^2 &= m_A^2\, s_\beta c_\beta, \\ \lambda_1 &= \frac{-s_\beta^2 m_A^2 + c_\alpha^2 m_H^2 + s_\alpha^2 m_h^2}{v^2 c_\beta^2}, \\ \lambda_2 &= \frac{-c_\beta^2 m_A^2 + s_\alpha^2 m_H^2 + c_\alpha^2 m_h^2}{v^2 s_\beta^2}, \\ \lambda_3 &= \frac{2m_{H^\pm}^2 - m_A^2}{v^2} + \frac{s_{2\alpha}(m_H^2 - m_h^2)}{v^2 s_{2\beta}}, \\ \lambda_4 &= \frac{2(m_A^2 - m_{H^\pm}^2)}{v^2}, \end{aligned} \tag{8}$$

where $c_\theta = \cos\theta$ and $s_\theta = \sin\theta$, for any needed angle θ, and the m's are the masses of the various scalars. The m_{11}^2 and m_{22}^2 parameters are obtained by solving the stationarity conditions $\partial V_H/(\partial v_1) = 0$ and $\partial V_H/(\partial v_2) = 0$. The angle α parametrizes the mixing

$$\begin{aligned} h &= \mathrm{Re}(\Phi_1^0)\, s_\alpha - \mathrm{Re}(\Phi_2^0)\, c_\alpha, \\ H &= -\mathrm{Re}(\Phi_1^0)\, c_\alpha - \mathrm{Re}(\Phi_2^0)\, s_\alpha, \end{aligned} \tag{9}$$

and can be taken between $-\pi/2$ and $\pi/2$. Eventually, the angles α and β will be determined by probing the couplings of the scalar particles to the fermions. Usually, two types of couplings to fermions are considered in theories with the $U(1)$ symmetry. In Type-I, all fermions couple to Φ_2; in Type-II, up-quarks couple to Φ_2, while down-quarks and charged leptons couple to Φ_1.

We have generated randomly 180.000 points by setting $v = 246\,\mathrm{GeV}$, $m_h = 125\,\mathrm{GeV}$, $m_H \in [125, 900]\,\mathrm{GeV}$, $m_A \in [10, 900]\,\mathrm{GeV}$, $m_{H^\pm} \in [90, 900]\,\mathrm{GeV}$, and $-\pi/2 \leq \alpha \leq \pi/2$. In addition, due to experimental bounds coming from B-physics, we have kept $\tan\beta > 1$. The corresponding couplings in Eqs. (8) have been forced to comply with a V_H bounded from below, to satisfy tree-level unitarity, and to be consistent with constraints from electroweak precision observables.

Fig. 1 shows the generated points in the (x, y) plane of Eqs. (4). In solid black lines we show the astroid. Points inside (outside) the astroid satisfy (do not satisfy) Eq. (3) and correspond to potentials which do (do not) have two minima. Green/light-gray points inside the astroid have $D > 0$, exhibiting two minima, with our current vacuum being the global minimum. This is a safe situation. Red/dark-gray points inside the astroid have $D < 0$, exhibit two minima, but correspond to the panic situation where our current vacuum lies above the other one. We have checked by explicit computation that the red/dark-gray points satisfy Eq. (6), while the green/light-gray points inside the astroid do not.



FIG. 1: Placement in the (x, y) plane of the points generated. Points inside the astroid (solid lines) have two minima; those in red/dark-gray correspond to the panic vacuum.

In order to study the panic vacuum in detail, we have generated two new sets of points where Eqs. (2)-(3) hold, and $D < 0$. The data sample for Type-I has 100.000 points. The Type-II model has the further constraint that $m_{H^\pm} > 340\,\mathrm{GeV}$, arising from $b \to s\gamma$ almost independently of $\tan\beta$ [10]. For Type-II, we have generated 58.000 points obeying this further constraint.

The generated points are used to calculate R_f, defined as the number of events predicted in the model for the process $pp \to h \to f$, divided by the prediction obtained in the SM for the same final state f. The experimental bounds at 1σ level are [9] $R_{ZZ} = 0.93 \pm 0.28$, $R_{\gamma\gamma} = 1.66 \pm 0.33$, summing over all production mechanisms.



FIG. 2: Predictions from Type-II in the $(R_{\gamma\gamma}, R_{ZZ})$ plane for panic vacuum points (in red/dark-gray) and for non-panic points (in green/light-gray). Also shown are the 1σ (solid line) and 2σ (dashed line) experimental bounds.



FIG. 3: Prediction for panic vacuum points in Type-II in the $(R_{\gamma\gamma}, R^{\mathrm{vbf}}_{\gamma\gamma})$ plane. They lie outside the 1σ (solid line) and 2σ (dashed line) ellipse bounds in Fig. 12 of Ref. [11].

The results we obtain for Type-II are shown in Figs. 2 and 3. Fig. 3 also includes the 1σ (solid line) and 2σ (dashed line) bounds coming from the ellipse in Fig. 12 of Ref. [11]. While Type-II (green/light-gray) points for which our current vacuum coincides with the global minimum are consistent with experiment, we find that Type-

II (red/dark-gray) panic points are excluded at the 2σ level by both ZZ and by $(R_{\gamma\gamma}, R_{\gamma\gamma}^{\rm vbf})$.

The results we obtain for Type-I are shown in the $(R_{\gamma\gamma}, R_{ZZ})$ plane of Fig. 4. We find that $R_{\gamma\gamma} \lesssim 0.88$, $R_{ZZ} \lesssim 1$, and the panic points lie outside the 2σ $R_{\gamma\gamma}$ experimental band. In contrast, the Type-I panic points we have generated in the $(R_{\gamma\gamma}, R_{\gamma\gamma}^{\rm vbf})$ plane lie outside the 1σ (elliptical) band, but inside the 2σ band. We could have $R_{\gamma\gamma}^{\rm vbf} > 1$, but only for values of $R_{\gamma\gamma} < 0.6$. Thus, we cannot reach the SM, $R_{\gamma\gamma} = R_{ZZ} = R_{\gamma\gamma}^{\rm vbf} = 1$, and the measurements of $R_{\gamma\gamma}$ are barely consistent with our results for Type-I.



FIG. 4: Predictions from Type-I in the $(R_{\gamma\gamma}, R_{ZZ})$ plane for panic vacuum points (in red/dark-gray) and for non-panic points (in green/light-gray). Also shown are the 1σ (solid line) and 2σ (dashed line) experimental bounds.

In summary, we have studied the vacuum structure of a 2HDM with a softly broken $U(1)$ symmetry, looking out for situations where our current vacuum has a larger energy density than the true global minimum of the scalar potential. We stress that the exclusion of panic vacuua should be addressed is models with extended Higgs sectors, where it should be implemented as an extra constraint on the theory's parameter space. Our conclusion is that, within our model, the recent LHC experiments are crucial in excluding the panic vacuum in Type-II and strongly disfavoring it in Type-I. If the current central values remain, the statistics accumulated by the end of this year will imply a precision that might also allow the exclusion of Type-I panic points. We can all sleep safer then... It is interesting that a cosmological question about the fate of our vacuum can be answered by laboratory experiments.

The works of A.B., P.M.F. and R.S. are supported in part by the Portuguese *Fundação para a Ciência e a Tecnologia* (FCT) under contract PTDC/FIS/117951/2010, by FP7 Reintegration Grant, number PERG08-GA-2010-277025, and by PEst-OE/FIS/UI0618/2011. The work of J.P.S. is funded by FCT through the projects CERN/FP/109305/2009 and U777-Plurianual, and by the EU RTN project Marie Curie: PITN-GA-2009-237920. I.P.I. is thankful to CFTC, University of Lisbon, for their hospitality. His work is supported by grants RFBR 11-02-00242-a, RF President grant for scientific schools NSc-3802.2012.2, and the Program of Department of Physics SC RAS and SB RAS "Studies of Higgs boson and exotic particles at LHC".

---

[*] E-mail: ferreira@cii.fc.ul.pt

[†] E-mail: igor.ivanov@ulg.ac.be

[‡] E-mail: rsantos@cii.fc.ul.pt

[§] E-mail: jpsilva@cftp.ist.utl.pt

[1] S. L. Glashow, Nucl. Phys. B **22**, 579 (1961). S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967). A. Salam, *Elementary Particles and Fields: Relativistic Groups and Analiticity (Nobel Symposium No. 8)*, edited by N. Svartholm (Almquist and Wiksell, Stockholm, 1968), p. 367.

[2] P. W. Higgs, Phys. Lett. **12**, 132 (1964). F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964). P. W. Higgs, Phys. Rev. Lett. **13**, 508 (1964). G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).

[3] For recent results, see G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]]. S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]].

[4] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38** (1977) 1440.

[5] For a recent review on the 2HDM, see, for example, G.C. Branco, P.M. Ferreira, L. Lavoura, M.N. Rebelo, M. Sher and J.P. Silva, Phys. Rept. **516**, 1 (2012) [arXiv:1106.0034 [hep-ph]].

[6] I. P. Ivanov, Phys. Rev. D **75**, 035001 (2007) [Erratum-ibid. D **76**, 039902 (2007)] [hep-ph/0609018]; *ibid*, **77**, 015017 (2008).

[7] N. Turok and J. Zadrozny, Nucl. Phys. B **369**, 729 (1992). I. P. Ivanov, Acta Phys. Polon. B **40**, 2789 (2009) [arXiv:0812.4984 [hep-ph]]. I. F. Ginzburg, I. P. Ivanov and K. A. Kanishev, Phys. Rev. D **81**, 085031 (2010) [arXiv:0911.2383 [hep-ph]]. I. F. Ginzburg, K. A. Kanishev, M. Krawczyk and D. Sokolowska, Phys. Rev. D **82**, 123533 (2010) [arXiv:1009.4593 [hep-ph]].

[8] See, for example, G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP **1208**, 098 (2012) [arXiv:1205.6497 [hep-ph]], and references therein.

[9] Since our main point does not depend on the specific details, we use the averages put forth in A. Arbey, M. Battaglia, A. Djouadi, F. Mahmoudi, arXiv:1211.4004, updated from *ibid.*, JHEP **1209** 107, (2012) [arXiv:1207.1348 [hep-ph]].

[10] F. Mahmoudi, talk given at Prospects For Charged Higgs Discovery At Colliders (CHARGED 2012), 8-11 October, Uppsala, Sweden.

[11] See Fig. 12 in ATLAS' article in Ref. [3].